

FIRST AID FOR SEIZURES

TONIC CLONIC SEIZURE

Person in Wheelchair



DO


- ✓ Stay with the person
- ✓ Time seizure
- ✓ Apply brakes
- ✓ Leave person in chair with seatbelt on
- ✓ Recline backrest only - do not tilt wheelchair
- ✓ Lean person slightly to one side to aid drainage of any fluid/food/vomit in mouth
- ✓ Support head and protect airway as required
- ✓ After the jerking stops carefully remove from chair and place in recovery position if possible or required
- ✓ Observe and reassure until recovered

DO NOT

- ✗ Put anything in the person's mouth
- ✗ Restrain the person
- ✗ Tilt or tip wheelchair

CALL 000 FOR AN AMBULANCE IF:

- You are in any doubt
- Injury has occurred
- There is food/fluid/vomit in mouth
- Seizure occurs in water
- Another seizure follows quickly
- Jerking of a tonic-clonic seizure lasts over 5 mins
- Complex partial seizures last longer than 15 mins
- Person has breathing difficulties after jerking stops



This is not medical advice or an exhaustive list of responses to seizures. This is a guide to help you consider your response to seizures. If you are in any doubt about what to do, do not hesitate to call an ambulance.

For more information or advice about epilepsy, contact Epilepsy Action on **1300 37 45 37** or visit **www.epilepsy.org.au**



EPILEPSY ACTION
A U S T R A L I A