

FIRST AID FOR SEIZURES

TONIC CLONIC SEIZURE

Person in Tilt-in-Space Wheelchair

Convulsive seizures where the body stiffens (tonic phase) followed by general muscle jerking (clonic phase).



DO


- ✓ Stay with the person
- ✓ Time seizure
- ✓ Apply brakes
- ✓ Leave person in chair with seatbelt on
- ✓ Tilt seat and lock in position
- ✓ Lean person slightly to one side to aid drainage of any food/fluid/vomit in mouth as required
- ✓ Support head and protect airway as required
- ✓ After the jerking stops carefully remove person from chair and place in recovery position as required
- ✓ Observe and reassure until recovered

DO NOT

- ✗ Put anything in the person's mouth
- ✗ Restrain the person
- ✗ Lift wheels from ground

CALL 000 FOR AN AMBULANCE IF:

- You are in any doubt
- Injury has occurred
- There is food/fluid/vomit in mouth
- Seizure occurs in water
- Person has breathing difficulties after jerking stops
- Another seizure quickly follows
- Seizure lasts longer than 5 mins
- The person is non-responsive for more than 5 mins after the seizure ends



This is not medical advice nor an exhaustive list of responses to seizures. This is a guide to help you consider your response to seizures. If you are in any doubt about what to do, do not hesitate to call an ambulance.



Epilepsy
Action
Australia

life changing impact